

Name _____

Date _____

* We solved for SIDES of triangles by using: \sin , \cos , and \tan . We will solve for ANGLES by using \sin^{-1} , \cos^{-1} , and \tan^{-1} .

$\sin^{-1}(x)$ is read as "the inverse sine of x ". Do not confuse $\sin^{-1}(x)$ as $\sin(x)$ having an exponent of -1.

1) Use a calculator to approximate the measures of the angles. Round your **final** answers to the nearest hundredth, if needed.

A) $\tan A = 0.75$

B) $\sin B = 0.87$






C) $\cos C = 0.15$

D) $\tan G = 0.43$


E) $\sin H = 0.68$

F) $\cos K = 0.94$

2) Approximate the measures of the angles. Round your **final** answers to the nearest hundredth, if needed.




3) Use your knowledge of Pythagorean Theorem, Triangle Sum Theorem, and Right Triangle Trigonometry to solve each triangle (find all missing angles and sides). Remember to keep all trig values to three decimals, minimum. Round your **final** answers to the nearest hundredth, if needed. Sides and angles opposite each other are commonly referred to using the same letter, with angles using a capital and side using a lowercase (i.e. opposite angle B, is side b).



A= _____ a= _____

B= _____ b= _____


C= _____ c= _____



A= _____ a= _____

B= _____ b= _____


C= _____ c= _____



A= _____ a= _____

B= _____ b= _____

C= _____ c= _____



A= _____ a= _____

B= _____ b= _____

C= _____ c= _____